



USDA, National Agricultural Statistics Service

Indiana Crop & Weather Report

Released: July 6, 2010

Vol. 60, WC070610

USDA, NASS, Indiana Field Office
1435 Win Hentschel Blvd.Suite 110
West Lafayette, IN 47906-4151(765) 494-8371
nass-in@nass.usda.gov

CROP REPORT FOR WEEK ENDING JULY 4

AGRICULTURAL SUMMARY

Dry weather returned to the state after several weeks of severe storms and rain, according to the Indiana Field Office of USDA's National Agricultural Statistics Service. Many fields with standing water had an opportunity to dry out. Some fields that had not been planted yet have now been planted with soybeans. Fields were sprayed this week that were previously too wet to access. Farmers used the break in the weather to continue harvesting wheat, with some reports of vomitoxin and low test weights. Double cropped soybeans were being planted after wheat harvest. Baling of hay and straw continued throughout the week.

FIELD CROPS REPORT

There were **5.3 days suitable for field work**. Thirty-one percent of the **corn** crop has **silked** compared with 2 percent last year and 8 percent for the 5-year average. Corn **condition** is rated 62 percent good to excellent compared with 64 percent last year at this time.

Ninety-six percent of the intended **soybean** acreage has **emerged** compared with 94 percent last year and 97 percent for the 5-year average. Twenty-three percent of the soybean acreage is **blooming** compared with 3 percent last year and 13 percent for the 5-year average. Soybean condition is rated 60 percent good to excellent compared with 64 percent last year.

Seventy-one percent of the **winter wheat** crop has been **harvested** compared with 52 percent last year and 51 percent for the 5-year average. Winter wheat **condition** is rated 57 percent good to excellent compared with 66 percent last year at this time.

Major activities during the week included: harvesting wheat, baling straw, spot planting soybeans in drowned out areas, herbicide applications, nitrogen applications to corn, mowing roadsides and ditches and taking care of livestock.

LIVESTOCK, PASTURE AND RANGE REPORT

Pasture condition is rated 76 percent good to excellent compared with 75 percent last year. Livestock are in good condition with ample pasture. The **first cutting of alfalfa hay** is 97 percent complete, while the **second cutting** is 23 percent complete.

CROP PROGRESS

| Crop | This Week | Last Week | Last Year | 5-Year Avg. |
|-------------------------|-----------|-----------|-----------|-------------|
| Percent | | | | |
| Corn Silked (Tasseled) | 31 | 8 | 2 | 8 |
| Soybeans Emerged | 96 | 90 | 94 | 97 |
| Soybeans Blooming | 23 | 7 | 3 | 13 |
| Winter Wheat Harvested | 71 | 37 | 52 | 51 |
| Alfalfa, First Cutting | 97 | 88 | 98 | 99 |
| Alfalfa, Second Cutting | 23 | NA | 26 | 32 |

CROP CONDITION

| Crop | Very Poor | Poor | Fair | Good | Excellent |
|--------------|-----------|------|------|------|-----------|
| Percent | | | | | |
| Corn | 3 | 9 | 26 | 46 | 16 |
| Soybean | 3 | 9 | 28 | 46 | 14 |
| Pasture | 1 | 4 | 19 | 53 | 23 |
| Winter Wheat | 1 | 8 | 34 | 49 | 8 |

SOIL MOISTURE & DAYS SUITABLE FOR FIELDWORK

| Soil Moisture | This Week | Last Week | Last Year |
|----------------------|-----------|-----------|-----------|
| Percent | | | |
| Topsoil | | | |
| Very Short | 1 | 1 | 1 |
| Short | 11 | 3 | 14 |
| Adequate | 66 | 43 | 67 |
| Surplus | 22 | 53 | 18 |
| Subsoil | | | |
| Very Short | 1 | 0 | 1 |
| Short | 6 | 2 | 7 |
| Adequate | 66 | 51 | 78 |
| Surplus | 27 | 47 | 14 |
| Days Suitable | 5.3 | 3.1 | 5.8 |

CONTACT INFORMATION



--Greg Preston, Director

--Michael Flanigan, Student Intern

E-mail Address: nass-in@nass.usda.gov

http://www.nass.usda.gov/Statistics_by_State/Indiana/

Crop Progress



Other Agricultural Comments And News

Tassel Emergence & Pollen Shed

Written by R.L. Nielsen, Purdue University

- Corn produces individual male and female flowers on the same plant.
- The tassel represents the male flower of the corn plant.

Depending on the year, Indiana's corn crop typically enters the critical flowering stages of pollen shed and silk emergence sometime between late June to late July. Success or failure during this period of the corn plant's life greatly influences the potential grain yield at harvest time.

As important as this process is to the determination of grain yield, it is surprising how little some folks know about the whole thing. Rather than leaving you to learn about such things "in the streets", I've developed this article and the accompanying one on silking ([Nielsen, 2010](#)) that describe the ins and outs of sex in the corn field.

Remember that corn has both male flowers and female flowers on the same plant (a flowering habit called [monoecious](#) for you trivia fans.) Interestingly, both flowers are initially bisexual (aka "perfect"), but during the course of development the female components (gynoecia) of the male flowers and the male components (stamens) of the female flowers abort, resulting in tassel (male) and ear (female) development.

Growth Stage VT (Tasseling)

Portions of the tassel may be visible before the plant technically reaches the last leaf stage (final visible leaf collar) has occurred. By definition, growth stage VT occurs when the last branch of the tassel emerges from the whorl ([Ritchie et al., 1993](#)). This authoritative source furthermore stated that growth stage VT is "initiated when the last branch of the tassel is completely visible and the silks have not yet emerged." Once upon a time, that exact developmental sequence may have been true, but today's hybrids tend to behave differently. It is not uncommon for silk emergence to begin not only prior to the last tassel branch appearing from the whorl, but also prior to the exertion of anthers and pollen shed ([Nielsen, 2009](#)).

Plant height reaches its maximum at or shortly after growth stage VT as the final stalk internodes complete their elongation. The corn plant is most vulnerable to hail damage at growth stage VT because all of its leaves are exposed. Complete (100 %) leaf loss at growth stage VT will usually result in complete (100 %) yield loss by harvest. Even if pollination results in successful fertilization of the ovules, entire ear shoots will usually die because so few leaves remain to produce the necessary carbohydrates (by photosynthesis) to complete grain fill.

Tassel Morphology

Approximately 1,000 individual spikelets form on each tassel and each one bears two florets encased in two large glumes. Each floret contains three anthers. An anther and its attached filament comprise the stamen of the male flower. The anthers are those "thingamajigs" that hang from the tassel during pollination. Under a magnifying lens, anthers look somewhat like the double barrel of a shotgun. Do the math and you will realize that an individual tassel produces approximately 6,000 pollen-bearing anthers, although hybrids can vary greatly for this number.

As these florets mature, elongation of the filaments helps exert the anthers from the glumes. Pollen is dispersed through pores that open at the tips of the anthers. Pollen shed usually begins in the mid-portion of the central tassel spike and then progresses upward, downward and outward over time. Anthers typically emerge from the upper floret of the pair first, while those from lower floret typically emerge later the same day or on following days. Spent anthers eventually drop from the tassel and are sometimes mistaken for the pollen when observed on the leaves or ground.

The yellow or white "dust-like" pollen that falls from a tassel represents millions of individual, nearly microscopic, spherical, yellowish- or whitish translucent pollen grains. Estimates of the total number of pollen grains produced per tassel range from 2 to 25 million. Each pollen grain contains the male genetic material necessary for fertilizing the ovary of one potential kernel.

(continued on back page)

Weather Information Table

Week Ending Sunday, July 4, 2010

| Station | Past Week Weather Summary Data | | | | | | | Accumulation | | | | |
|--------------------------|--------------------------------|----|-----|---------|-------|------|------|-----------------------|-------|------|---------------|------|
| | Air | | | | | | Avg | April 1, 2010 through | | | | |
| | Temperature | | | Precip. | | | 4 in | July 4, 2010 | | | | |
| | | | | | | | Soil | Precipitation | | | GDD Base 50°F | |
| | Hi | Lo | Avg | DFN | Total | Days | Temp | Total | DFN | Days | Total | DFN |
| Northwest (1) | | | | | | | | | | | | |
| Chalmers_5W | 87 | 50 | 68 | -7 | 1.18 | 1 | | 20.76 | +9.01 | 42 | 1342 | +114 |
| Francesville | 86 | 47 | 68 | -5 | 0.00 | 0 | | 13.58 | +1.66 | 38 | 1325 | +215 |
| Valparaiso_AP_I | 86 | 49 | 69 | -4 | 0.24 | 1 | | 14.91 | +2.31 | 41 | 1327 | +253 |
| Wanatah | 86 | 45 | 68 | -5 | 0.35 | 1 | 75 | 14.78 | +2.82 | 36 | 1238 | +220 |
| Winamac | 87 | 50 | 69 | -4 | 0.50 | 1 | | 16.89 | +4.97 | 44 | 1371 | +261 |
| North Central (2) | | | | | | | | | | | | |
| Plymouth | 86 | 47 | 67 | -7 | 0.35 | 1 | | 13.52 | +1.10 | 33 | 1263 | +104 |
| South_Bend | 86 | 48 | 68 | -4 | 0.26 | 1 | | 14.13 | +2.46 | 38 | 1305 | +250 |
| Young_America | 88 | 42 | 68 | -6 | 0.22 | 1 | | 20.12 | +8.66 | 36 | 1358 | +233 |
| Northeast (3) | | | | | | | | | | | | |
| Fort_Wayne | 90 | 53 | 71 | -3 | 0.55 | 1 | | 15.54 | +4.65 | 40 | 1523 | +412 |
| Kendallville | 87 | 49 | 68 | -5 | 0.59 | 1 | | 13.29 | +1.83 | 46 | 1252 | +206 |
| West Central (4) | | | | | | | | | | | | |
| Greencastle | 87 | 51 | 68 | -8 | 1.32 | 1 | | 17.38 | +4.54 | 41 | 1372 | +65 |
| Perrysville | 89 | 54 | 71 | -4 | 0.09 | 1 | 84 | 15.22 | +2.53 | 38 | 1597 | +386 |
| Spencer_Ag | 89 | 53 | 70 | -5 | 1.70 | 1 | | 21.77 | +8.34 | 41 | 1499 | +291 |
| Terre_Haute_AFB | 90 | 50 | 71 | -4 | 1.01 | 1 | | 17.96 | +5.44 | 44 | 1662 | +361 |
| W_Lafayette_6NW | 88 | 48 | 69 | -4 | 0.23 | 1 | 82 | 17.52 | +5.73 | 34 | 1476 | +344 |
| Central (5) | | | | | | | | | | | | |
| Eagle_Creek_AP | 88 | 56 | 72 | -4 | 1.54 | 2 | | 16.23 | +4.50 | 40 | 1682 | +394 |
| Greenfield | 90 | 53 | 71 | -4 | 1.59 | 1 | | 21.28 | +8.82 | 42 | 1538 | +323 |
| Indianapolis_AP | 89 | 56 | 73 | -3 | 0.70 | 1 | | 17.13 | +5.40 | 37 | 1734 | +446 |
| Indianapolis_SE | 87 | 51 | 69 | -6 | 0.84 | 1 | | 18.48 | +6.52 | 38 | 1493 | +230 |
| Tipton_Ag | 92 | 49 | 69 | -4 | 1.04 | 1 | 77 | 17.55 | +5.82 | 42 | 1409 | +322 |
| East Central (6) | | | | | | | | | | | | |
| Farmland | 89 | 50 | 68 | -4 | 1.06 | 2 | 77 | 16.91 | +4.91 | 47 | 1422 | +374 |
| New_Castle | 88 | 49 | 67 | -6 | 0.86 | 2 | | 20.23 | +7.24 | 40 | 1358 | +282 |
| Southwest (7) | | | | | | | | | | | | |
| Evansville | 94 | 56 | 75 | -3 | 0.60 | 1 | | 8.83 | -3.93 | 33 | 1948 | +397 |
| Freelandville | 92 | 57 | 73 | -3 | 1.51 | 1 | | 16.48 | +3.35 | 36 | 1733 | +377 |
| Shoals_8S | 91 | 51 | 70 | -6 | 1.16 | 1 | | 17.19 | +3.18 | 28 | 1569 | +272 |
| Stendal | 94 | 57 | 75 | -2 | 0.91 | 1 | | 11.93 | -2.38 | 30 | 1978 | +540 |
| Vincennes_5NE | 92 | 58 | 73 | -3 | 0.73 | 1 | 83 | 14.57 | +1.44 | 38 | 1778 | +422 |
| South Central (8) | | | | | | | | | | | | |
| Leavenworth | 92 | 58 | 74 | -1 | 0.92 | 2 | | 13.92 | -0.26 | 47 | 1759 | +461 |
| Oolitic | 87 | 54 | 71 | -4 | 2.45 | 1 | 82 | 20.30 | +6.96 | 39 | 1539 | +313 |
| Tell_City | 94 | 58 | 75 | -3 | 0.41 | 1 | | 13.27 | -1.03 | 27 | 1887 | +433 |
| Southeast (9) | | | | | | | | | | | | |
| Brookville | 90 | 51 | 71 | -3 | 0.73 | 1 | | 15.96 | +3.34 | 38 | 1541 | +408 |
| Greensburg | 90 | 55 | 72 | -2 | 0.53 | 1 | | 17.77 | +4.81 | 38 | 1695 | +492 |
| Seymour | 88 | 53 | 70 | -5 | 2.77 | 1 | | 16.50 | +3.96 | 33 | 1534 | +288 |

Copyright 2010: Agricultural Weather Information Service, Inc.
All rights reserved.

DFN = Departure From Normal.
GDD = Growing Degree Days.
Precipitation (Rainfall or melted snow/ice) in inches.
Precipitation Days = Days with precip of .01 inch or more.
Air Temperatures in Degrees Fahrenheit.

For more weather information, visit www.awis.com
or call 1-888-798-9955.

Tassel Emergence & Pollen Shed (continued)

The outer membrane of a pollen grain is very thin. Once dispersed into the atmosphere, pollen grains remain viable for only a few minutes before they desiccate. Yet, with only a 15 mph wind, pollen grains can travel as far as 1/2 mile within those couple of minutes.

Therein lies the concern of the potential for pollen “drift” from a transgenic corn field to an adjacent non-transgenic corn field and the risk of transgenic “contamination” of grain intended for non-transgenic markets. The good news is that recent research suggests that the overwhelming majority of a corn field’s pollen load is shed in the field itself.

All of the pollen from a single anther may be released in as little as three minutes. All the anthers on an individual tassel may take as long as seven days to finish shedding pollen, although the greatest volume of pollen is typically shed during the second and third day of anther emergence. Because of natural field variability in plant development, a whole field may take as long as 14 days to complete pollen shed.

Peak pollen shed usually occurs in mid-morning. Some research indicates that pollen shed decreases after

temperatures surpass 86F. A second “flush” of pollen often occurs in late afternoon or evening as temperatures cool. Pollen shed may occur throughout most of the day under relatively cool, cloudy conditions.

Weather conditions influence pollen shed. If the anthers are wet, the pores will not open and pollen will not be released. Thus, on an average Indiana summer morning following a heavy evening dew, pollen shed will not begin until the dew dries and the anther pores open. Similarly, pollen is not shed during rainy conditions. Cool, humid temperatures delay pollen shed, while hot, dry conditions hasten pollen shed.

Extreme heat stress (100F or greater) can kill corn pollen, but fortunately the plant avoids significant pollen loss by virtue of two developmental characteristics. First of all, corn pollen does not mature or shed all at once. Pollen maturity and shed occur over several days and up to two weeks. Therefore, a day or two of extreme heat usually does not affect the entire pollen supply. More importantly, the majority of daily pollen shed occurs in the morning hours when air temperature is much more moderate.

The INDIANA CROP & WEATHER REPORT (USPS 675-770), (ISSN43-817X) is issued weekly April through November by the USDA, NASS Indiana Field Office, 1435 Win Hentschel Blvd, Suite 110, West Lafayette, IN 47906-4151. For information on subscribing, send request to above address. POSTMASTER: Send address change to the USDA, NASS, Indiana Field Office, 1435 Win Hentschel Blvd, Suite 110, West Lafayette, IN 47906-4151.

WEEKLY NEWS REPORT

FIRST-CLASS MAIL
POSTAGE & FEES PAID
USDA
PERMIT NO. G-38

INDIANA CROP & WEATHER REPORT
USDA, NASS, INDIANA FIELD OFFICE
1435 WIN HENTSCHEL BLVD, STE 110
WEST LAFAYETTE IN 47906-4151